

University of Groningen

The Westerbork Hydrogen Accretion in Local Galaxies (HALOGAS) survey (Corrigendum)

Heald, G.; Józsa, G.; Serra, P.; Zschaechner, L.; Rand, R.; Fraternali, F.; Oosterloo, T.; Walterbos, R.; Jütte, E.; Gentile, G.

Published in:
Astronomy & astrophysics

DOI:
[10.1051/0004-6361/201015938e](https://doi.org/10.1051/0004-6361/201015938e)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2012

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Heald, G., Józsa, G., Serra, P., Zschaechner, L., Rand, R., Fraternali, F., Oosterloo, T., Walterbos, R., Jütte, E., & Gentile, G. (2012). The Westerbork Hydrogen Accretion in Local Galaxies (HALOGAS) survey (Corrigendum): I. Survey description and pilot observations. *Astronomy & astrophysics*, 544, [C1].
<https://doi.org/10.1051/0004-6361/201015938e>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

The Westerbork Hydrogen Accretion in Local GALaxies (HALOGAS) survey

I. Survey description and pilot observations (Corrigendum)

G. Heald¹, G. Józsa¹, P. Serra¹, L. Zschaechner², R. Rand², F. Fraternali³, T. Oosterloo^{1,4}, R. Walterbos⁵,
 E. Jütte⁶, and G. Gentile⁷

- ¹ Netherlands Institute for Radio Astronomy (ASTRON), Postbus 2, 7990 AA Dwingeloo, The Netherlands
² University of New Mexico, 800 Yale Blvd, Albuquerque, NM, USA
³ Astronomy Department, Bologna University, via Ranzani 1, 40127 Bologna, Italy
⁴ Kapteyn Astronomical Institute, Postbus 800, 9700 AV Groningen, The Netherlands
⁵ Department of Astronomy, New Mexico State University, PO Box 30001, MSC 4500, Las Cruces, NM 88003, USA
⁶ Astronomisches Institut der Ruhr-Universität Bochum, Universitätsstr. 150, 44780 Bochum, Germany
⁷ Sterrenkundig Observatorium, Ghent University, Krijgslaan 281, S9, 9000 Ghent, Belgium

A&A 526, A118 (2011), DOI: [10.1051/0004-6361/201015938](https://doi.org/10.1051/0004-6361/201015938)

Key words. galaxies: spiral – galaxies: evolution – galaxies: ISM – galaxies: halos – galaxies: kinematics and dynamics – errata, addenda




Fig. 1. Comparison between corrected SFR values and the original values. The line represents equal SFR values. Most corrected SFR values are higher, because d_{T88} tends to be lower than our adopted distances.

We discovered that in the original paper, the adopted distances (d_{best}) were not used in the calculation of the star formation rate (SFR) given in Col. (12) of Table 1. Instead we inadvertently used the Tully (1988) distances (d_{T88}). Here we correct this error, for reference in future HALOGAS publications. The newly computed SFRs are compared to the originally published values in Fig. 1. The results of the original paper are not affected by this correction. Our adopted distances are unchanged from the original paper, but are repeated here for clarity. A description of the origin of the adopted distances can be found in the original paper. A description of the methods used for calculating the SFR values themselves can also be found in the original paper.

References

- Moshir, M., et al. 1990, in IRAS Faint Source Catalogue, version 2.0
 Tully, R. B. 1988, Nearby galaxies catalog (Cambridge Univ. Press)

Table 1. Updated SFR values for the HALOGAS sample.

UGC	Other IDs	d_{T88} (Mpc)	d_{best} (Mpc)	Old SFR ($M_{\odot} \text{ yr}^{-1}$)	New SFR ($M_{\odot} \text{ yr}^{-1}$)
1256	NGC 0672	7.5	7.6	0.22	0.23
1831	NGC 0891	9.6	9.2	2.4	2.2
1913	NGC 0925	9.4	9.1	0.83	0.77
1983	NGC 0949	10.3	11.3	0.26	0.31
2082	–	10.7	14.4	0.023	0.041
2137	NGC 1003	10.7	11.6	0.34	0.40
3918	NGC 2403	4.2	3.2	1.0	0.60
4278	IC 2233	10.6	13.6	0.11	0.18
4284	NGC 2541	10.6	12.0	<0.27 ^a	<0.35 ^a
5572	NGC 3198	10.8	14.5	0.61	1.1
7045	NGC 4062	9.7	16.9	0.22	0.67
7322	NGC 4244	3.1	4.4	0.058	0.12
7353	NGC 4258	6.8	7.6	1.4	1.7
	(M 106)				
7377	NGC 4274	9.7	19.4	0.31	1.2
7539	NGC 4414	9.7	17.8	1.3	4.2
7591	NGC 4448	9.7	9.7	0.056	0.056
7766	NGC 4559	9.7	7.9	1.1	0.69
7772	NGC 4565	9.7	10.8	0.54	0.67
7774	–	6.8	24.4	0.0074	0.095
7865	NGC 4631	6.9	7.6	2.1	2.6
8286	NGC 5023	6.0	6.6	0.032	0.039
8334	NGC 5055	7.2	8.5	1.5	2.1
	(M 63)				
8550	NGC 5229	6.4	5.1	0.014	0.0086
9179	NGC 5585	7.0	8.7	0.26	0.41

Notes. ^(a) The SFR value for NGC 2541 is strictly speaking an upper limit, because the IRAS 25 μm flux is catalogued as a non-detection by Moshir et al. (1990).